







Optical Gel Can Be Used Almost Anywhere





Installs Easily in Minutes

Protects: Ledges, Signs, Balconies, Soffits, Beams, Skylights, Signs, Roofs, AC Equipment, Enclosed Spaces


Optical Gel Installation Guidelines

1. Clean site thoroughly (see below)
2. For pigeons, place dishes as shown below. For other birds, see below.

Visit www.birdbarrier.com for cleaning solutions to sanitize the site prior to Optical Gel™ installation.

- **Dissolve-It** » for breaking down bird droppings and eliminating odors
- **MDBC-10** » a safe, concentrated formula for cleaning droppings


Nesting Sites: 6 in. maximum spacing



Night Roosting Sites: 8 in. maximum spacing



Day Roosting Sites: 10 in. maximum spacing



DIRECTIONS FOR USE

Bird Barrier Optical Gel™ may be used to deter feral pigeons, gulls, and many other unwanted birds from roosting on buildings and other structures. Bird Barrier Optical Gel™ is supplied in ready-to-use plastic-dish containers. These should be affixed to surfaces with commercially available silicone adhesive or other suitable and removable adhesive. (See **CLEANING SURFACES PRIOR TO TREATMENT** and **APPLICATION DIRECTIONS** below.)

APPLICATION DIRECTIONS

Remove covers from dishes and affix dishes to dry surfaces with silicone or other removable adhesive.

For feral pigeons, dishes should be spaced (center-to-center) as follows:

- Nesting sites: 6 inches
- Night roosts: 8 inches
- Day roosts: 10 inches

General guidelines for spacing of dishes for other species of bird are:

- Gull: 12 inches
- Corvids (crow, magpie, etc): 10 inches
- Starling: 8 inches.

Some testing may be needed to determine best placement for other bird species (see **INSTALLATION INSTRUCTIONS** at www.birdbarrier.com).

STORAGE AND DISPOSAL

Nonrefillable container. Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in a cool, dry place inaccessible to children and pets. Do not store with solvents that may dissolve the plastic dishes.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of at an approved waste disposal facility. Cover exposed gel in dishes to prevent spreading prior to disposal.

Container Handling: Nonrefillable containers. Do not reuse or refill containers. Offer for recycling, if available.

CLEANING STRUCTURE SURFACES PRIOR TO TREATMENT PRECAUTIONARY STATEMENTS *(skip this step if structure is already clean)*

Wear protective eyewear and face mask or respirator to avoid inhaling bird feces, nesting materials, feathers, and dead birds, while cleaning structure surfaces before placing Bird Barrier Optical Gel™ containers. Wash skin and clothing after cleaning bird debris and before eating or drinking. Do not touch or rub eyes while or after handling Bird Barrier Optical Gel™.

For Bird Barrier Optical Gel™ to be most effective, bird nesting materials, feces, feathers, and dead birds must be removed prior to application of the Optical Gel™ containers. To reduce potential exposure to diseases from bird debris, persons that clean or remove bird debris, should wear waterproof gloves and appropriate personal protective equipment (PPE) for dust and aerosol mist protection while cleaning areas to be treated. Thoroughly clean surfaces, including confined spaces such as building attics and other protected areas, where birds roost or nest.

Take precautions to ensure nearby people are not exposed to debris from cleaning.

The following PPE should be used during the cleaning process when cleaning may result in exposure, especially respiratory exposure:

- Long-sleeved shirt and long pants. Wash thoroughly after use or wear disposable outer clothing, such as Tyvek® disposable coveralls or suits.
- Waterproof gloves or disposable waterproof gloves.
- Rubber boots or other waterproof disposable foot protection.

To avoid respiratory exposure, especially in confined areas, cleaners should wear a dust/mist filtering respirator meeting NIOSH standards of at least N-95, R-95, or P-95. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions are available for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Use a biocide or water mixed with detergent to wet surfaces where feces are present to limit generation of airborne dust and aerosol mists during cleaning. Allow cleaned surfaces to dry prior to treatment. When cleaning large amounts of bird debris, workers should be informed of the possible health risks involved. Persons with compromised immune systems should avoid the cleaning operation.

If pressure-washers are used to strip off dried feces, dust and aerosol control measures, such as containing the area with plastic sheeting, should be taken. Wetting down the work area will prevent inhalation, reduce the risk of disease from bird feces and debris, and will also help prevent the spread of dust and aerosols.

An alternative to use of a pressure washer is to soak the feces and debris with water and then use a shovel to remove them. The wet material should be collected in heavy-duty plastic trash bags or some other type of secure container and discarded with trash.

For technical support or other questions, contact Bird Barrier at 310-527-8000.